PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Charles-LeMoyne Date: 2025-08-15 23:46

Anesthésiste / Anesthetist: Dr. Patricia Wong

Chirurgien / Surgeon: Dr. Sophie Chen Assistant(s): Dr. resident John Paul

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH INTUSSUSCEPTION.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH INTUSSUSCEPTION.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH OMENTAL WRAPPING.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and inflamed tissue

Anesthésie / Anesthesia: General anesthesia with nitrous oxide

Historique et constatations opératoires / History and operative findings:

Pediatric patient (11, male) presenting with acute onset abdominal pain with rebound tenderness. History: no prior abdominal surgery. Imaging confirmed appendicitis.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. Total intravenous anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. A vertical infraumbilical incision is made and carried down to the fascia which is incised sharply. Three trocars in total are used for laparoscopic access. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, extensive adhesions noted. The appendix was phlegmonous with moderate inflammation. Multiple small abscesses were encountered. No significant adhesions. Dissection is carried out to isolate the base of the appendix. The mesoappendix is dissected and the appendiceal artery is controlled with electrocautery. The appendix is ligated at its base with two absorbable sutures and then amputated. Specimen placed in EndoCatch bag for removal. Abdominal lavage performed until clear. The umbilical fascia is reapproximated with interrupted Maxon 2-0 sutures. Skin incisions are closed with Dermabond. No evidence of peritoneal carcinomatosis.

The patient will be continued on IV antibiotics and advanced to diet as tolerated. Incidental Meckel's diverticulum found and left in situ. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-JWDSQF-11161