PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Jewish General Hospital Date: 2025-08-03 11:25

Anesthésiste / Anesthetist: Dr. Rachel Stein

Chirurgien / Surgeon: Dr. Olivia Davis Assistant(s): Dr. resident Carlos Mendez

Diagnostic préopératoire / Pre-operative diagnosis:

LOCALIZED PERITONITIS SECONDARY TO APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

LOCALIZED PERITONITIS SECONDARY TO APPENDICITIS.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH DRAINAGE OF ABSCESS.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and peri-appendiceal tissue

Anesthésie / Anesthesia: General anesthesia with caudal block

Historique et constatations opératoires / History and operative findings:

2-year-old female with 3 days abdominal pain. Treated for Crohn's disease; symptoms persisted. Imaging: MRI demonstrating RLQ inflammation. Past medical history is otherwise unremarkable.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. A small infraumbilical incision is made and the abdominal cavity is entered under direct vision. Additional 5 mm trocars are placed in the right and left lower quadrants under laparoscopic guidance. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, moderate adhesions noted. Findings include ruptured appendix with severe inflammatory changes. There was a contained abscess in the right lower quadrant. Minimal adhesions were noted. Patient tolerated procedure well. Meticulous dissection performed due to distorted anatomy. The appendiceal artery is ligated and divided. The base of the appendix is healthy and we place three EndoLoops - two proximal and one distal - before transecting the appendix. Specimen placed in EndoCatch bag for removal. Abdominal lavage performed until clear. All port sites closed with Dermabond.

We will continue current antibiotic regimen and begin enteral feeds when bowel function returns. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-DXX7QP-13879