PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: St. Mary's Hospital Date: 2025-07-31 14:14

Anesthésiste / Anesthetist: Dr. Paul Anderson Chirurgien / Surgeon: Dr. Robert Tremblay Assistant(s): Dr. resident Marc Gagnon

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH FREE FLUID.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH FREE FLUID.

Opération / Operation:

OPEN APPENDECTOMY.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and peri-appendiceal tissue

Anesthésie / Anesthesia: General anesthesia and epidural block

Historique et constatations opératoires / History and operative findings:

A 8-year-old male with 1 day history of abdominal pain and constipation. Failed conservative management for urinary tract infection. Imaging: ultrasound showing appendicitis. Past medical history is otherwise unremarkable.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with mask induction administered. Time-out was performed and abdomen prepped in sterile fashion. We create a 1 cm infraumbilical incision and enter the abdomen using the Hasson technique. Single-incision laparoscopic port is used. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, extensive adhesions noted. Operative findings included suppurative appendix and fluctuating inflammation. Purulent fluid was noted throughout the abdominal cavity. Minimal adhesions were noted. The appendix is mobilized using a combination of sharp and blunt dissection. Mesenteric vessels to the appendix are secured prior to removal. The appendix is ligated at its base with two absorbable sutures and then amputated. Specimen placed in EndoCatch bag for removal. We irrigate the abdominal cavity copiously with warm saline until the effluent is clear. Fascial closure is performed at the umbilical site using Vicryl 3-0. The skin is closed with Dermabond.

Monitor for signs of infection; advance diet as tolerated. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-FBEQWU-12761