# PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital de Verdun Date: 2025-03-19 13:25

Anesthésiste / Anesthetist: Dr. Claire Dubois Chirurgien / Surgeon: Dr. Robert Tremblay Assistant(s): Dr. resident Emily Clark

Diagnostic préopératoire / Pre-operative diagnosis:

# APPENDICITIS WITH ABSCESS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH ABSCESS.

Opération / Operation:

### LAPAROSCOPIC APPENDECTOMY WITH OMENTAL WRAPPING.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with surrounding lymph nodes

Anesthésie / Anesthesia: General anesthesia with mask induction

#### Historique et constatations opératoires / History and operative findings:

A 15-year-old male with 1 day history of abdominal pain with elevated WBC. Failed conservative management for constipation. Imaging: ultrasound suggestive of appendicitis.

# Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. Direct trocar insertion after skin and fascia incision. Two working ports are established in the right and left lower quadrants. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, moderate adhesions noted. Intraoperative examination revealed necrotic appendix. Multiple small abscesses were encountered. Dense adhesions were encountered during dissection. The surrounding tissues showed diffuse reaction. Bladder and ureters visualized, no injury. Meticulous dissection performed due to distorted anatomy. The mesoappendix is dissected and the appendiceal artery is controlled with electrocautery. Endoscopic stapling is used for appendiceal division. Specimen placed in EndoCatch bag for removal. We irrigate the abdominal cavity copiously with warm saline until the effluent is clear. Fascial closure is performed at the umbilical site using PDS 3-0. The skin is closed with subcuticular Vicryl 4-0. Small serosal tear repaired intraoperatively.

Monitor wound sites for infection. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-570NHG-13910