PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Royal Victoria Hospital Date: 2024-09-30 07:58

Anesthésiste / Anesthetist: Dr. Thomas White

Chirurgien / Surgeon: Dr. Victor Chen Assistant(s): Dr. resident Chloe Nguyen

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH SEPSIS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH MESENTERIC LYMPHADENITIS.

Opération / Operation:

OPEN APPENDECTOMY WITH PELVIC LAVAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and inflamed tissue

Anesthésie / Anesthesia: General endotracheal anesthesia

Historique et constatations opératoires / History and operative findings:

1-year-old male with several hours abdominal pain, markedly elevated WBC, high CRP, high fever. Imaging: MRI showing bowel wall thickening, family history of appendicitis.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. Total intravenous anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. A vertical infraumbilical incision is made and carried down to the fascia which is incised sharply. Two working ports are established in the right and left lower quadrants. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, extensive adhesions noted. Appendix appeared shrunken, surrounded by extensive inflammatory reaction. There was a contained abscess in the right lower quadrant. Minimal adhesions were noted. We carefully dissect the inflammatory mass and identify the necrotic appendix. The appendiceal mesentery is carefully taken down with harmonic scalpel. Endoscopic stapling is used for appendiceal division. Specimen placed in EndoCatch bag for removal. Saline is used for thorough irrigation of all quadrants. All port sites closed with Steri-Strips.

Consult infectious disease if antibiotics need adjustment. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-QUSIME-11380