PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Maisonneuve-Rosemont Date: 2025-09-16 03:54

Anesthésiste / Anesthetist: Dr. Claire Dubois Chirurgien / Surgeon: Dr. Marie-Claire Dubois

Assistant(s): Dr. resident Jake Turner

Diagnostic préopératoire / Pre-operative diagnosis:

COMPLICATED APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH PERITONEAL CONTAMINATION.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH OMENTAL WRAPPING.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and mesoappendix

Anesthésie / Anesthesia: General anesthesia with caudal block

Historique et constatations opératoires / History and operative findings:

Patient (17 years, non-binary) presented with abdominal pain with fever, markedly elevated WBC, high CRP. Imaging: ultrasound showing phlegmonous appendicitis. Past medical history is otherwise unremarkable.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia and epidural block administered. Time-out was performed and abdomen prepped in sterile fashion. A small infraumbilical incision is made and the abdominal cavity is entered under direct vision. Accessory port placed in epigastric region. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, extensive adhesions noted. Appendix appeared gangrenous, surrounded by localized inflammatory reaction. No pus or abscess formation found. Mild adhesions between bowel loops observed. Minor bleeding controlled with cautery. Surrounding omentum and bowel are separated from the inflammatory mass. The appendiceal mesentery is carefully taken down with harmonic scalpel. The appendix is ligated at its base with two absorbable sutures and then amputated. Specimen placed in EndoCatch bag for removal. We irrigate the abdominal cavity copiously with warm saline until the effluent is clear. We close the fascia at the umbilical port with figure-of-eight sutures of PDS 2-0 and the skin with interrupted silk 4-0. Patient tolerated procedure well.

Repeat CBC and CRP postoperatively. Ovaries and uterus normal in female patients. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-RSPBV7-13229