## PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Royal Victoria Hospital Date: 2024-06-30 21:32

Anesthésiste / Anesthetist: Dr. Camille Roy Chirurgien / Surgeon: Dr. Sophie Chen Assistant(s): Dr. resident Ethan Wright

Diagnostic préopératoire / Pre-operative diagnosis:

## **RUPTURED APPENDICITIS.**

Diagnostic postopératoire / Post-operative diagnosis:

PERFORATED APPENDICITIS WITH ABSCESS.

Opération / Operation:

#### LAPAROSCOPIC APPENDECTOMY.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with surrounding lymph nodes Anesthésie / Anesthesia: General anesthesia with local infiltration

# Historique et constatations opératoires / History and operative findings:

5-year-old male with one week abdominal pain. Treated for renal colic; symptoms persisted. Imaging: CT scan confirming appendicitis.

# Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with local infiltration administered. Time-out was performed and abdomen prepped in sterile fashion. Transverse infraumbilical incision is performed and access gained via blunt dissection. Three trocars in total are used for laparoscopic access. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, extensive adhesions noted. The surgical field demonstrated hyperemic appendix with localized surrounding inflammation. No pus or abscess formation found. No abnormal adhesions found. We carefully dissect the inflammatory mass and identify the friable appendix. Mesenteric vessels to the appendix are secured prior to removal. Endoscopic stapling is used for appendiceal division. Specimen placed in EndoCatch bag for removal. Copious irrigation is undertaken to ensure removal of all inflammatory debris. The umbilical fascia is reapproximated with interrupted Vicryl 3-0 sutures. Skin incisions are closed with Steri-Strips. No evidence of peritoneal carcinomatosis.

The patient will be continued on IV antibiotics and advanced to diet as tolerated. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-5D2E7W-11873