PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Ste-Agathe Hospital Date: 2025-07-26 03:50

Anesthésiste / Anesthetist: Dr. Amélie Moreau

Chirurgien / Surgeon: Dr. James Wilson Assistant(s): Dr. resident Zoe Tremblay

Diagnostic préopératoire / Pre-operative diagnosis:

PERFORATED APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

ACUTE APPENDICITIS.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH DRAINAGE OF ABSCESS.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with attached omentum

Anesthésie / Anesthesia: General anesthesia with mask induction

Historique et constatations opératoires / History and operative findings:

A 14-year-old non-binary who presented with abdominal pain with vomiting. Initially evaluated 2 days prior and diagnosed with renal colic. Now has markedly elevated WBC, high CRP, high fever. Imaging: CT scan confirming appendicitis.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with nitrous oxide administered. Time-out was performed and abdomen prepped in sterile fashion. A small infraumbilical incision is made and the abdominal cavity is entered under direct vision. Single-incision laparoscopic port is used. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, minimal adhesions noted. Operative findings included friable appendix and minimal inflammation. No abscess was identified. The omentum was wrapped around the inflamed appendix. Dissection is carried out to isolate the base of the appendix. We dissect the mesentery of the appendix and use the electrocautery to coagulate the artery. We secure the appendiceal base with two Endoloops and transect between them. Specimen placed in EndoCatch bag for removal. Saline is used for thorough irrigation of all quadrants. Fascial closure is performed at the umbilical site using PDS 2-0. The skin is closed with subcuticular Vicryl 4-0.

Monitor wound sites for infection. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-92H0ML-12272