## PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: CHU Sainte-Justine Date: 2025-01-23 15:15

Anesthésiste / Anesthetist: Dr. Claire Dubois Chirurgien / Surgeon: Dr. James Wilson Assistant(s): Dr. resident Carlos Mendez

Diagnostic préopératoire / Pre-operative diagnosis:

## APPENDICITIS WITH BOWEL OBSTRUCTION.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH BOWEL OBSTRUCTION.

Opération / Operation:

#### OPEN APPENDECTOMY WITH PELVIC LAVAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and inflamed tissue

Anesthésie / Anesthesia: General endotracheal anesthesia

# Historique et constatations opératoires / History and operative findings:

8-year-old female with 1 day abdominal pain. Treated for constipation; symptoms persisted. Imaging: ultrasound suggestive of appendicitis.

# Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General endotracheal anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. We make an infraumbilical incision of 1 cm, dissect the subcutaneous tissue bluntly and penetrate the abdominal cavity via an open technique. Additional 5 mm trocars are placed in the right and left lower quadrants under laparoscopic guidance. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, diffuse adhesions noted. Appendix appeared thick-walled, surrounded by extensive inflammatory reaction. Abscess cavity found in RLQ and irrigated. Minimal adhesions were noted. We carefully dissect the inflammatory mass and identify the phlegmonous appendix. We dissect the mesentery of the appendix and use the electrocautery to coagulate the artery. We secure the appendiceal base with two Endoloops and transect between them. Specimen placed in EndoCatch bag for removal. Irrigation performed with antibiotic solution. The umbilical fascia is reapproximated with interrupted Polysorb 2-0 sutures. Skin incisions are closed with Steri-Strips.

Consult infectious disease if antibiotics need adjustment. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-KMUS1E-13878