# PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Pierre-Boucher Date: 2025-08-22 09:14

Anesthésiste / Anesthetist: Dr. Patricia Wong

Chirurgien / Surgeon: Dr. Paul Lambert Assistant(s): Dr. resident Jake Turner

Diagnostic préopératoire / Pre-operative diagnosis:

### **RUPTURED APPENDICITIS.**

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH MESENTERIC LYMPHADENITIS.

Opération / Operation:

#### OPEN APPENDECTOMY WITH PELVIC LAVAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix Anesthésie / Anesthesia: General anesthesia with regional block

### Historique et constatations opératoires / History and operative findings:

A 15-year-old female with 1 day history of abdominal pain after trauma. Failed conservative management for food poisoning. Imaging: CT scan showing peri-appendiceal fluid. Recent travel history may be relevant.

## Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. Direct trocar insertion after skin and fascia incision. Additional 5 mm trocars are placed in the right and left lower quadrants under laparoscopic guidance. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, moderate adhesions noted. Findings include necrotic appendix with extensive inflammatory changes. Abscess cavity found in RLQ and irrigated. No significant adhesions. Minimal intraoperative blood loss. Dissection is carried out to isolate the base of the appendix. The mesoappendix is dissected and the appendiceal artery is controlled with electrocautery. We secure the appendiceal base with two Endoloops and transect between them. Specimen placed in EndoCatch bag for removal. Saline is used for thorough irrigation of all quadrants. Umbilical port site is closed with Vicryl 3-0 and skin with subcuticular Monocryl 4-0.

The patient will be continued on IV antibiotics and advanced to diet as tolerated. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-CPXGIH-12607