## PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: CHU Sainte-Justine Date: 2024-06-06 21:09

Anesthésiste / Anesthetist: Dr. Rachel Stein

Chirurgien / Surgeon: Dr. Victor Chen Assistant(s): Dr. resident Zoe Tremblay

Diagnostic préopératoire / Pre-operative diagnosis:

## LOCALIZED PERITONITIS SECONDARY TO APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

SUPPURATIVE APPENDICITIS.

Opération / Operation:

#### LAPAROSCOPIC APPENDECTOMY.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and peri-appendiceal tissue

Anesthésie / Anesthesia: General anesthesia with local infiltration

# Historique et constatations opératoires / History and operative findings:

A 8-year-old male with one week history of abdominal pain with fever. Failed conservative management for renal colic. Imaging: ultrasound showing appendicitis.

# Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. Total intravenous anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. Transverse infraumbilical incision is performed and access gained via blunt dissection. Two working ports are established in the right and left lower quadrants. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, minimal adhesions noted. The surgical field demonstrated friable appendix with mild surrounding inflammation. No pus or abscess formation found. The appendix was adhered to surrounding structures. Appendix is isolated after adhesiolysis. The mesoappendix is divided using a bipolar energy device. We secure the appendiceal base with two Endoloops and transect between them. Specimen placed in EndoCatch bag for removal. Saline is used for thorough irrigation of all quadrants. Umbilical port site is closed with Ethibond 2-0 and skin with interrupted silk 4-0.

IV fluids and pain management as per protocol. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-NK1ZVY-10639