## PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: St. Mary's Hospital Date: 2025-08-07 10:25

Anesthésiste / Anesthetist: Dr. Lisa Garcia Chirurgien / Surgeon: Dr. Elena Rodriguez

Assistant(s): Dr. resident Leo Morel

Diagnostic préopératoire / Pre-operative diagnosis:

# APPENDICITIS WITH FREE FLUID.

Diagnostic postopératoire / Post-operative diagnosis:

**ACUTE APPENDICITIS.** 

Opération / Operation:

### OPEN APPENDECTOMY WITH PELVIC LAVAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with attached omentum

Anesthésie / Anesthesia: General anesthesia with local infiltration

#### Historique et constatations opératoires / History and operative findings:

Pediatric patient (9, non-binary) presenting with acute onset RLQ tenderness and guarding. History: previous similar episode. Imaging confirmed appendicitis. Past medical history is otherwise unremarkable.

# Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with regional block administered. Time-out was performed and abdomen prepped in sterile fashion. We create a 1 cm infraumbilical incision and enter the abdomen using the Hasson technique. Two working ports are established in the right and left lower quadrants. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, significant adhesions noted. Appendix was friable and surrounded by No pus or abscess formation found. and The appendix was adhered to surrounding structures. We carefully dissect the inflammatory mass and identify the acutely inflamed appendix. The mesoappendix is divided using a bipolar energy device. The appendix is ligated at its base with two absorbable sutures and then amputated. Specimen placed in EndoCatch bag for removal. Abdominal lavage performed until clear. We close the fascia with Vicryl 2-0 in a interrupted fashion. Skin is approximated with interrupted nylon 4-0.

Continue surgical care with antibiotic therapy and diet advancement as tolerated. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-8VF710-10910