PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: CHU Sainte-Justine Date: 2024-10-12 22:47

Anesthésiste / Anesthetist: Dr. Julia Miller Chirurgien / Surgeon: Dr. Elena Rodriguez

Assistant(s): Dr. resident Leo Morel

Diagnostic préopératoire / Pre-operative diagnosis:

SUPPURATIVE APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH EXTENSIVE ADHESIONS.

Opération / Operation:

APPENDECTOMY WITH INTRA-ABDOMINAL DRAINAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and inflamed tissue

Anesthésie / Anesthesia: General anesthesia with regional block

Historique et constatations opératoires / History and operative findings:

Pediatric patient (2, male) presenting with acute onset abdominal pain and constipation. History: history of constipation. Imaging confirmed appendicitis.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. Total intravenous anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. We make an infraumbilical incision of 1 cm, dissect the subcutaneous tissue bluntly and penetrate the abdominal cavity via an open technique. Two working ports are established in the right and left lower quadrants. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, diffuse adhesions noted. Appendix appeared sclerotic, surrounded by persistent inflammatory reaction. No abscess, but turbid fluid present. Minimal adhesions were noted. Minor bleeding controlled with cautery. We proceed with careful dissection of the appendiceal attachments. The appendiceal mesentery is carefully taken down with harmonic scalpel. The base of the appendix is healthy and we place three EndoLoops - two proximal and one distal - before transecting the appendix. Specimen placed in EndoCatch bag for removal. Abdominal lavage performed until clear. All port sites closed with Dermabond.

Pain control with acetaminophen and morphine as needed. No need for drains postoperatively. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-60CQ07-13562