## PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Shriners Hospitals for Children Date: 2024-07-28 09:36

Anesthésiste / Anesthetist: Dr. Omar Fahmy

Chirurgien / Surgeon: Dr. Paul Lambert Assistant(s): Dr. resident Chloe Nguyen

Diagnostic préopératoire / Pre-operative diagnosis:

## APPENDICITIS WITH INTUSSUSCEPTION.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH MESENTERIC LYMPHADENITIS.

Opération / Operation:

### LAPAROSCOPIC APPENDECTOMY.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and mesoappendix

Anesthésie / Anesthesia: General anesthesia with nitrous oxide

### Historique et constatations opératoires / History and operative findings:

10-year-old female with 3 days abdominal pain. Treated for constipation; symptoms persisted. Imaging: ultrasound reporting lymphadenopathy. Recent travel history may be relevant.

# Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with local infiltration administered. Time-out was performed and abdomen prepped in sterile fashion. A vertical infraumbilical incision is made and carried down to the fascia which is incised sharply. Additional 5 mm trocars are placed in the right and left lower quadrants under laparoscopic guidance. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, minimal adhesions noted. The appendix was hyperemic with minimal inflammation. A small localized abscess was found and drained. The appendix was adhered to surrounding structures. Appendix is isolated after adhesiolysis. The appendiceal artery is ligated and divided. Endoscopic stapling is used for appendiceal division. Specimen placed in EndoCatch bag for removal. Copious irrigation is undertaken to ensure removal of all inflammatory debris. We close the fascia at the umbilical port with figure-of-eight sutures of Polysorb 2-0 and the skin with Steri-Strips.

Discharge home when tolerating oral intake and afebrile. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-MND642-10150