PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: CHU de Québec Date: 2025-04-06 21:31

Anesthésiste / Anesthetist: Dr. John Evans

Chirurgien / Surgeon: Dr. Victor Chen Assistant(s): Dr. resident Zoe Tremblay

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH ABSCESS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH ABSCESS.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH OMENTAL WRAPPING.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix Anesthésie / Anesthesia: General anesthesia with nitrous oxide

Historique et constatations opératoires / History and operative findings:

A 1-year-old non-binary with several hours history of abdominal pain with palpable mass. Failed conservative management for Crohn's disease. Imaging: ultrasound with non-visualized appendix and secondary signs. Past medical history is otherwise unremarkable.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with caudal block administered. Time-out was performed and abdomen prepped in sterile fashion. Incision is made in left lower quadrant for open conversion. Additional 5 mm trocars are placed in the right and left lower quadrants under laparoscopic guidance. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, significant adhesions noted. Appendix appeared acutely inflamed, surrounded by persistent inflammatory reaction. There was a contained abscess in the right lower quadrant. No abnormal adhesions found. No evidence of peritoneal carcinomatosis. The appendix is mobilized using a combination of sharp and blunt dissection. We dissect the mesentery of the appendix and use the electrocautery to coagulate the artery. Endoscopic stapling is used for appendiceal division. Specimen placed in EndoCatch bag for removal. Irrigation performed with antibiotic solution. The umbilical fascia is reapproximated with interrupted Ethibond 2-0 sutures. Skin incisions are closed with interrupted nylon 4-0.

Repeat CBC and CRP postoperatively. No technical difficulties encountered during surgery. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-0LII49-12955