PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Pierre-Boucher Date: 2024-11-06 12:32

Anesthésiste / Anesthetist: Dr. Rachel Stein Chirurgien / Surgeon: Dr. James Wilson Assistant(s): Dr. resident Marc Gagnon

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH ABSCESS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH EXTENSIVE ADHESIONS.

Opération / Operation:

LAPAROSCOPIC CONVERTED TO OPEN APPENDECTOMY.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with attached omentum

Anesthésie / Anesthesia: Total intravenous anesthesia

Historique et constatations opératoires / History and operative findings:

A 1-year-old male who presented with abdominal pain with diarrhea. Initially evaluated 3 days prior and diagnosed with viral syndrome. Now has markedly elevated WBC, elevated CRP, high fever. Imaging: ultrasound showing appendicitis.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with nitrous oxide administered. Time-out was performed and abdomen prepped in sterile fashion. Incision is made in left lower quadrant for open conversion. Three trocars in total are used for laparoscopic access. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, minimal adhesions noted. Appendix appeared hyperemic, surrounded by extensive inflammatory reaction. A large pelvic abscess was present and evacuated. No abnormal adhesions found. Surrounding omentum and bowel are separated from the inflammatory mass. We dissect the mesentery of the appendix and use the electrocautery to coagulate the artery. The base of the appendix is healthy and we place three EndoLoops - two proximal and one distal - before transecting the appendix. Specimen placed in EndoCatch bag for removal. Saline is used for thorough irrigation of all quadrants. All port sites closed with interrupted nylon 4-0.

Consult infectious disease if antibiotics need adjustment. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-O4NL9K-12919