PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: St. Mary's Hospital Date: 2025-02-08 13:09

Anesthésiste / Anesthetist: Dr. David Smith

Chirurgien / Surgeon: Dr. Aisha Patel Assistant(s): Dr. resident Lucas Martin

Diagnostic préopératoire / Pre-operative diagnosis:

ACUTE APPENDICITIS WITH PERITONITIS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH MESENTERIC LYMPHADENITIS.

Opération / Operation:

OPEN APPENDECTOMY.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with surrounding lymph nodes

Anesthésie / Anesthesia: General anesthesia with nitrous oxide

Historique et constatations opératoires / History and operative findings:

A 17-year-old female who presented with persistent vomiting and abdominal pain. Initially evaluated 2 days prior and diagnosed with IBD. Now has markedly elevated WBC, high CRP, no fever. Imaging: CT scan revealing free fluid.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with regional block administered. Time-out was performed and abdomen prepped in sterile fashion. Incision is made in left lower quadrant for open conversion. We place two additional trocars, one in the suprapubic region and one in the left lower quadrant, both under direct vision. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, diffuse adhesions noted. Findings include phlegmonous appendix with marked inflammatory changes. Abscess cavity found in RLQ and irrigated. No significant adhesions. Appendix is isolated after adhesiolysis. Mesenteric vessels to the appendix are secured prior to removal. Endoscopic stapling is used for appendiceal division. Specimen placed in EndoCatch bag for removal. Copious irrigation is undertaken to ensure removal of all inflammatory debris. Fascial closure is performed at the umbilical site using PDS 3-0. The skin is closed with interrupted nylon 4-0.

Postoperative imaging if fever persists. No intraoperative complications occurred. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-2XIOYV-11817