PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: CHU de Québec Date: 2024-05-03 07:37

Anesthésiste / Anesthetist: Dr. David Smith

Chirurgien / Surgeon: Dr. Samuel Lee Assistant(s): Dr. resident Chloe Nguyen

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH INTUSSUSCEPTION.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH PELVIC ABSCESS.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH OMENTAL WRAPPING.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendicolith Anesthésie / Anesthesia: General anesthesia with regional block

Historique et constatations opératoires / History and operative findings:

17-year-old male with one week abdominal pain. Treated for intussusception; symptoms persisted. Imaging: CT scan showing appendiceal abscess.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with regional block administered. Time-out was performed and abdomen prepped in sterile fashion. We make an infraumbilical incision of 1 cm, dissect the subcutaneous tissue bluntly and penetrate the abdominal cavity via an open technique. Single-incision laparoscopic port is used. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, significant adhesions noted. The appendix was hyperemic with fluctuating inflammation. A large pelvic abscess was present and evacuated. Dense adhesions were encountered during dissection. Appendix is isolated after adhesiolysis. The mesoappendix is dissected and the appendiceal artery is controlled with electrocautery. Endoscopic stapling is used for appendiceal division. Specimen placed in EndoCatch bag for removal. Abdominal lavage performed until clear. All port sites closed with interrupted silk 4-0. Minimal intraoperative blood loss.

The patient will be continued on IV antibiotics and advanced to diet as tolerated. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-9WSIVU-11698