PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Ste-Agathe Hospital Date: 2024-01-04 02:24

Anesthésiste / Anesthetist: Dr. Patricia Wong

Chirurgien / Surgeon: Dr. Aisha Patel Assistant(s): Dr. resident Jennifer Park

Diagnostic préopératoire / Pre-operative diagnosis:

LOCALIZED PERITONITIS SECONDARY TO APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH MESENTERIC LYMPHADENITIS.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH DRAINAGE OF ABSCESS.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and peri-appendiceal tissue

Anesthésie / Anesthesia: General anesthesia with local infiltration

Historique et constatations opératoires / History and operative findings:

Pediatric patient (4, female) presenting with acute onset abdominal pain and constipation. History: recent antibiotic use. Imaging confirmed appendicitis.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. Direct trocar insertion after skin and fascia incision. Three trocars in total are used for laparoscopic access. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, minimal adhesions noted. The surgical field demonstrated necrotic appendix with mild surrounding inflammation. No pus or abscess formation found. No significant adhesions. No intraoperative complications occurred. Appendix is isolated after adhesiolysis. We dissect the mesentery of the appendix and use the electrocautery to coagulate the artery. We secure the appendiceal base with two Endoloops and transect between them. Specimen placed in EndoCatch bag for removal. Saline is used for thorough irrigation of all quadrants. We close the fascia with Vicryl 2-0 in a interrupted fashion. Skin is approximated with subcuticular Monocryl 4-0. No technical difficulties encountered during surgery.

The patient will be continued on IV antibiotics and advanced to diet as tolerated. No evidence of peritoneal carcinomatosis. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-JMTA2U-13322