PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital de Verdun Date: 2024-03-05 05:09

Anesthésiste / Anesthetist: Dr. Amélie Moreau

Chirurgien / Surgeon: Dr. Samuel Lee Assistant(s): Dr. resident Lucas Martin

Diagnostic préopératoire / Pre-operative diagnosis:

COMPLICATED APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH EXTENSIVE ADHESIONS.

Opération / Operation:

LAPAROSCOPIC CONVERTED TO OPEN APPENDECTOMY.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with surrounding lymph nodes

Anesthésie / Anesthesia: General anesthesia with local infiltration

Historique et constatations opératoires / History and operative findings:

5-year-old male with 3 days abdominal pain, elevated WBC, normal CRP, high fever. Imaging: MRI showing bowel wall thickening. history of constipation. Past medical history is otherwise unremarkable.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with caudal block administered. Time-out was performed and abdomen prepped in sterile fashion. We create a 1 cm infraumbilical incision and enter the abdomen using the Hasson technique. Three trocars in total are used for laparoscopic access. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, diffuse adhesions noted. The surgical field demonstrated phlegmonous appendix with fluctuating surrounding inflammation. There was a contained abscess in the right lower quadrant. Mild adhesions between bowel loops observed. We carefully dissect the inflammatory mass and identify the gangrenous appendix. The appendiceal mesentery is carefully taken down with harmonic scalpel. Base of appendix secured with purse-string suture prior to removal. Specimen placed in EndoCatch bag for removal. Thorough irrigation of the abdominal cavity is performed, removing all purulent material. Umbilical port site is closed with Maxon 2-0 and skin with non-absorbable Prolene 4-0.

Monitor for signs of infection; advance diet as tolerated. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-P4LKFJ-11269