PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: CHU Sainte-Justine Date: 2025-09-26 14:26

Anesthésiste / Anesthetist: Dr. Omar Fahmy Chirurgien / Surgeon: Dr. Martin Levesque Assistant(s): Dr. resident Carlos Mendez

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS MIMICKING OVARIAN PATHOLOGY.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS MIMICKING OVARIAN PATHOLOGY.

Opération / Operation:

APPENDECTOMY WITH INTRA-ABDOMINAL DRAINAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and abscess wall

Anesthésie / Anesthesia: General anesthesia with caudal block

Historique et constatations opératoires / History and operative findings:

1-year-old female with one week abdominal pain. Treated for gastroenteritis; symptoms persisted. Imaging: ultrasound suggestive of appendicitis. Past medical history is otherwise unremarkable.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with caudal block administered. Time-out was performed and abdomen prepped in sterile fashion. A vertical infraumbilical incision is made and carried down to the fascia which is incised sharply. Supplementary ports are placed in the suprapubic area and left iliac fossa. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, moderate adhesions noted. The surgical field demonstrated friable appendix with marked surrounding inflammation. No abscess, but turbid fluid present. Minimal adhesions were noted. We proceed with careful dissection of the appendiceal attachments. The mesoappendix is dissected and the appendiceal artery is controlled with electrocautery. The appendix is ligated at its base with two absorbable sutures and then amputated. Specimen placed in EndoCatch bag for removal. Saline is used for thorough irrigation of all quadrants. The umbilical fascia is reapproximated with interrupted Polysorb 2-0 sutures. Skin incisions are closed with subcuticular Monocryl 4-0.

Monitor wound sites for infection. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-UINIS3-14037