## PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: CHU Sainte-Justine Date: 2024-07-08 04:03

Anesthésiste / Anesthetist: Dr. Lisa Garcia Chirurgien / Surgeon: Dr. Isabelle Girard Assistant(s): Dr. resident Sophia Lee

Diagnostic préopératoire / Pre-operative diagnosis:

## APPENDICITIS WITH SEPSIS.

Diagnostic postopératoire / Post-operative diagnosis:

**ACUTE APPENDICITIS.** 

Opération / Operation:

### LAPAROSCOPIC CONVERTED TO OPEN APPENDECTOMY.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with attached omentum

Anesthésie / Anesthesia: General anesthesia with nitrous oxide

### Historique et constatations opératoires / History and operative findings:

Patient (14 years, male) presented with right lower quadrant pain, markedly elevated WBC, high CRP. Imaging: CT scan showing appendiceal abscess.

# Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. Incision is made in left lower quadrant for open conversion. We place two additional trocars, one in the suprapubic region and one in the left lower quadrant, both under direct vision. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, diffuse adhesions noted. Appendix was suppurative and surrounded by Abscess cavity found in RLQ and irrigated. and Minimal adhesions were noted. Surrounding omentum and bowel are separated from the inflammatory mass. The mesoappendix is dissected and the appendiceal artery is controlled with electrocautery. Endoscopic stapling is used for appendiceal division. Specimen placed in EndoCatch bag for removal. Copious irrigation is undertaken to ensure removal of all inflammatory debris. We close the fascia at the umbilical port with figure-of-eight sutures of Vicryl 3-0 and the skin with subcuticular Monocryl 4-0.

Patient to receive postoperative IV antibiotics with monitoring for return of bowel function. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-1YPJC5-13723