## PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Pierre-Boucher Date: 2025-07-20 16:14

Anesthésiste / Anesthetist: Dr. Claire Dubois

Chirurgien / Surgeon: Dr. Samuel Lee Assistant(s): Dr. resident Fatima Sheikh

Diagnostic préopératoire / Pre-operative diagnosis:

## PHLEGMONOUS APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

**GANGRENOUS APPENDICITIS.** 

Opération / Operation:

#### LAPAROSCOPIC APPENDECTOMY WITH OMENTAL WRAPPING.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with surrounding lymph nodes

Anesthésie / Anesthesia: General anesthesia and epidural block

# Historique et constatations opératoires / History and operative findings:

9-year-old female with 1 day abdominal pain. Treated for IBD; symptoms persisted. Imaging: CT scan showing appendiceal abscess.

## Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with mask induction administered. Time-out was performed and abdomen prepped in sterile fashion. We make an infraumbilical incision of 1 cm, dissect the subcutaneous tissue bluntly and penetrate the abdominal cavity via an open technique. Additional 5 mm trocars are placed in the right and left lower quadrants under laparoscopic guidance. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, minimal adhesions noted. The appendix was acutely inflamed with patchy inflammation. Purulent fluid was noted throughout the abdominal cavity. Fibrinous adhesions were lysed during the procedure. Appendix is isolated after adhesiolysis. Appendiceal vessels controlled with clips. We secure the appendiceal base with two Endoloops and transect between them. Specimen placed in EndoCatch bag for removal. Thorough irrigation of the abdominal cavity is performed, removing all purulent material. We close the fascia at the umbilical port with figure-of-eight sutures of PDS 2-0 and the skin with Dermabond.

Patient to receive postoperative IV antibiotics with monitoring for return of bowel function. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-5LBPDD-12827