

PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Pierre-Boucher

Date: 2025-01-30 00:06

Anesthésiste / Anesthetist: Dr. David Smith

Chirurgien / Surgeon: Dr. Isabelle Girard

Assistant(s): Dr. resident Sophia Lee

Diagnostic préopératoire / Pre-operative diagnosis:

LOCALIZED PERITONITIS SECONDARY TO APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH PELVIC ABSCESS.

Opération / Operation:

OPEN APPENDECTOMY WITH PELVIC LAVAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendicolith

Anesthésie / Anesthesia: General endotracheal anesthesia

Historique et constatations opératoires / History and operative findings:

5-year-old non-binary with 3 days abdominal pain, normal WBC, normal CRP, low-grade fever. Imaging: ultrasound showing perforated appendicitis. recent antibiotic use.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. We create a 1 cm infraumbilical incision and enter the abdomen using the Hasson technique. Accessory port placed in epigastric region. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, significant adhesions noted. Intraoperative examination revealed perforated appendix. There was a contained abscess in the right lower quadrant. Minimal adhesions were noted. The surrounding tissues showed minimal reaction. Blunt dissection is used to free the appendix from surrounding structures. The mesoappendix is dissected and the appendiceal artery is controlled with electrocautery. The base of the appendix is healthy and we place three EndoLoops - two proximal and one distal - before transecting the appendix. Specimen placed in EndoCatch bag for removal. Thorough irrigation of the abdominal cavity is performed, removing all purulent material. We close the fascia with Vicryl 3-0 in an interrupted fashion. Skin is approximated with Dermabond. No evidence of peritoneal carcinomatosis.

Early ambulation and supportive care recommended. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-ST4RHS-10082