PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: CHU Sainte-Justine Date: 2024-03-07 10:37

Anesthésiste / Anesthetist: Dr. Rachel Stein Chirurgien / Surgeon: Dr. Sophie Chen

Assistant(s): Dr. resident Chloe Nguyen

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH ABSCESS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH HEMORRHAGE.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH IRRIGATION AND DRAINAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and inflamed tissue

Anesthésie / Anesthesia: General anesthesia with nitrous oxide

Historique et constatations opératoires / History and operative findings:

Patient (4 years, male) presented with abdominal pain with rebound tenderness, normal WBC, elevated CRP. Imaging: CT scan confirming appendicitis.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with mask induction administered. Time-out was performed and abdomen prepped in sterile fashion. Transverse infraumbilical incision is performed and access gained via blunt dissection. Supplementary ports are placed in the suprapubic area and left iliac fossa. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, minimal adhesions noted. Operative findings included gangrenous appendix and moderate inflammation. No abscess, but turbid fluid present. Mild adhesions between bowel loops observed. The appendix is mobilized using a combination of sharp and blunt dissection. We dissect the mesentery of the appendix and use the electrocautery to coagulate the artery. Base of appendix secured with purse-string suture prior to removal. Specimen placed in EndoCatch bag for removal. Thorough irrigation of the abdominal cavity is performed, removing all purulent material. The umbilical fascia is reapproximated with interrupted Polysorb 2-0 sutures. Skin incisions are closed with interrupted silk 4-0.

Continue surgical care with antibiotic therapy and diet advancement as tolerated. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-80G09X-12861