PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: CHU de Québec Date: 2025-03-04 21:39

Anesthésiste / Anesthetist: Dr. Michael Brown Chirurgien / Surgeon: Dr. Sarah Johnson Assistant(s): Dr. resident Zoe Tremblay

Diagnostic préopératoire / Pre-operative diagnosis:

ACUTE APPENDICITIS WITH PERITONITIS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH HEMORRHAGE.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH DRAINAGE OF ABSCESS.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and abscess wall

Anesthésie / Anesthesia: General anesthesia with nitrous oxide

Historique et constatations opératoires / History and operative findings:

Pediatric patient (12, non-binary) presenting with acute onset abdominal pain with vomiting. History: family history of appendicitis. Imaging confirmed appendicitis. Recent travel history may be relevant.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia and epidural block administered. Time-out was performed and abdomen prepped in sterile fashion. We make an infraumbilical incision of 1 cm, dissect the subcutaneous tissue bluntly and penetrate the abdominal cavity via an open technique. Supplementary ports are placed in the suprapubic area and left iliac fossa. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, diffuse adhesions noted. The appendix was perforated with patchy inflammation. There was a contained abscess in the right lower quadrant. Minimal adhesions were noted. Appendix is isolated after adhesiolysis. The appendiceal artery is ligated and divided. Absorbable ligatures are applied prior to amputation. Specimen placed in EndoCatch bag for removal. We irrigate the abdominal cavity copiously with warm saline until the effluent is clear. We close the fascia at the umbilical port with figure-of-eight sutures of Ethibond 2-0 and the skin with interrupted nylon 4-0.

Monitor for signs of infection; advance diet as tolerated. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-5YKO8I-11477