PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Pierre-Boucher Date: 2025-06-13 03:33

Anesthésiste / Anesthetist: Dr. Paul Anderson

Chirurgien / Surgeon: Dr. Paul Lambert Assistant(s): Dr. resident Maya Singh

Diagnostic préopératoire / Pre-operative diagnosis:

SUPPURATIVE APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

SUPPURATIVE APPENDICITIS.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH OMENTAL WRAPPING.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix Anesthésie / Anesthesia: General anesthesia with regional block

Historique et constatations opératoires / History and operative findings:

12-year-old non-binary with several hours abdominal pain, elevated WBC, elevated CRP, high fever. Imaging: ultrasound showing perforated appendicitis. family history of appendicitis.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with mask induction administered. Time-out was performed and abdomen prepped in sterile fashion. A vertical infraumbilical incision is made and carried down to the fascia which is incised sharply. Supplementary ports are placed in the suprapubic area and left iliac fossa. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, minimal adhesions noted. Operative findings included thick-walled appendix and persistent inflammation. Multiple small abscesses were encountered. No significant adhesions. No bowel injury noted. We proceed with careful dissection of the appendiceal attachments. Appendiceal vessels controlled with clips. Endoscopic stapling is used for appendiceal division. Specimen placed in EndoCatch bag for removal. Thorough irrigation of the abdominal cavity is performed, removing all purulent material. We close the fascia at the umbilical port with figure-of-eight sutures of Vicryl 3-0 and the skin with subcuticular Monocryl 4-0.

Patient to receive postoperative IV antibiotics with monitoring for return of bowel function. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-7NJYWF-13303