PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital de Verdun Date: 2024-11-14 07:36

Anesthésiste / Anesthetist: Dr. Omar Fahmy Chirurgien / Surgeon: Dr. Robert Tremblay Assistant(s): Dr. resident Jake Turner

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH FREE FLUID.

Diagnostic postopératoire / Post-operative diagnosis:

ACUTE APPENDICITIS.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH OMENTAL WRAPPING.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with attached omentum

Anesthésie / Anesthesia: General anesthesia with caudal block

Historique et constatations opératoires / History and operative findings:

2-year-old male with 2 days abdominal pain. Treated for urinary tract infection; symptoms persisted. Imaging: MRI showing bowel wall thickening.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General endotracheal anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. We make an infraumbilical incision of 1 cm, dissect the subcutaneous tissue bluntly and penetrate the abdominal cavity via an open technique. Supplementary ports are placed in the suprapubic area and left iliac fossa. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, moderate adhesions noted. Findings include acutely inflamed appendix with severe inflammatory changes. No pus or abscess formation found. Mild adhesions between bowel loops observed. Appendix is isolated after adhesiolysis. We dissect the mesentery of the appendix and use the electrocautery to coagulate the artery. Base of appendix secured with purse-string suture prior to removal. Specimen placed in EndoCatch bag for removal. We lavage the abdomen extensively, paying particular attention to the pelvis and right gutter. The umbilical fascia is reapproximated with interrupted Maxon 2-0 sutures. Skin incisions are closed with interrupted nylon 4-0.

Pain control with acetaminophen and morphine as needed. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-7FI8A7-13529