

PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Montreal Children's

Date: 2024-12-17 06:01

Anesthésiste / Anesthetist: Dr. Rachel Stein

Chirurgien / Surgeon: Dr. Paul Lambert

Assistant(s): Dr. resident Chloe Nguyen

Diagnostic préopératoire / Pre-operative diagnosis:

LOCALIZED PERITONITIS SECONDARY TO APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

LOCALIZED PERITONITIS SECONDARY TO APPENDICITIS.

Opération / Operation:

APPENDECTOMY WITH REMOVAL OF APPENDICOLITH.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with attached omentum

Anesthésie / Anesthesia: General anesthesia with nitrous oxide

Historique et constatations opératoires / History and operative findings:

17-year-old non-binary with 3 days abdominal pain, elevated WBC, elevated CRP, high fever. Imaging: MRI demonstrating RLQ inflammation. no prior abdominal surgery.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with local infiltration administered. Time-out was performed and abdomen prepped in sterile fashion. We create a 1 cm infraumbilical incision and enter the abdomen using the Hasson technique. Three trocars in total are used for laparoscopic access. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, moderate adhesions noted. The surgical field demonstrated necrotic appendix with severe surrounding inflammation. A small localized abscess was found and drained. Minimal adhesions were noted. No bowel injury noted. We carefully dissect the inflammatory mass and identify the gangrenous appendix. The mesoappendix is divided using a bipolar energy device. We apply an endoscopic stapler to the base of the appendix and divide it. Specimen placed in EndoCatch bag for removal. Irrigation performed with antibiotic solution. Fascial closure is performed at the umbilical site using PDS 3-0. The skin is closed with Steri-Strips. No need for drains postoperatively.

We will continue current antibiotic regimen and begin enteral feeds when bowel function returns. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-SS7YGA-13985