## PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: MCH Date: 2024-09-28 18:21

Anesthésiste / Anesthetist: Dr. Claire Dubois Chirurgien / Surgeon: Dr. Martin Levesque Assistant(s): Dr. resident Jennifer Park

Diagnostic préopératoire / Pre-operative diagnosis:

## SUPPURATIVE APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

SUPPURATIVE APPENDICITIS.

Opération / Operation:

#### OPEN APPENDECTOMY.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and inflamed tissue

Anesthésie / Anesthesia: General anesthesia with mask induction

# Historique et constatations opératoires / History and operative findings:

A 10-year-old male who presented with abdominal pain with lethargy. Initially evaluated 1 days prior and diagnosed with urinary tract infection. Now has elevated WBC, high CRP, no fever. Imaging: CT scan confirming appendicitis. Recent travel history may be relevant.

## Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with local infiltration administered. Time-out was performed and abdomen prepped in sterile fashion. A small infraumbilical incision is made and the abdominal cavity is entered under direct vision. Supplementary ports are placed in the suprapubic area and left iliac fossa. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, significant adhesions noted. Appendix appeared necrotic, surrounded by severe inflammatory reaction. A large pelvic abscess was present and evacuated. No significant adhesions. Meticulous dissection performed due to distorted anatomy. The appendiceal artery is ligated and divided. We apply an endoscopic stapler to the base of the appendix and divide it. Specimen placed in EndoCatch bag for removal. We irrigate the abdominal cavity copiously with warm saline until the effluent is clear. We close the fascia at the umbilical port with figure-of-eight sutures of Vicryl 3-0 and the skin with interrupted nylon 4-0.

Postoperative imaging if fever persists. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-BWVGCQ-13658