PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Notre-Dame Date: 2024-03-13 09:53

Anesthésiste / Anesthetist: Dr. David Smith Chirurgien / Surgeon: Dr. Elena Rodriguez Assistant(s): Dr. resident Marc Gagnon

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH FREE FLUID.

Diagnostic postopératoire / Post-operative diagnosis:

PERFORATED APPENDICITIS.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH IRRIGATION AND DRAINAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and mesoappendix

Anesthésie / Anesthesia: General anesthesia with local infiltration

Historique et constatations opératoires / History and operative findings:

Patient (17 years, male) presented with abdominal pain with anorexia, normal WBC, high CRP. Imaging: ultrasound suggestive of appendicitis.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. We make an infraumbilical incision of 1 cm, dissect the subcutaneous tissue bluntly and penetrate the abdominal cavity via an open technique. We place two additional trocars, one in the suprapubic region and one in the left lower quadrant, both under direct vision. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, minimal adhesions noted. Findings include perforated appendix with localized inflammatory changes. A large pelvic abscess was present and evacuated. Dense adhesions were encountered during dissection. We proceed with careful dissection of the appendiceal attachments. The appendiceal mesentery is carefully taken down with harmonic scalpel. Base of appendix secured with purse-string suture prior to removal. Specimen placed in EndoCatch bag for removal. Thorough irrigation of the abdominal cavity is performed, removing all purulent material. Fascial closure is performed at the umbilical site using Vicryl 2-0. The skin is closed with subcuticular Monocryl 4-0. No technical difficulties encountered during surgery.

Monitor wound sites for infection. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-Y2B5D3-12791