PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Pierre-Boucher Date: 2025-03-17 12:12

Anesthésiste / Anesthetist: Dr. Claire Dubois

Chirurgien / Surgeon: Dr. Paul Lambert Assistant(s): Dr. resident Marc Gagnon

Diagnostic préopératoire / Pre-operative diagnosis:

PERFORATED APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

ACUTE APPENDICITIS.

Opération / Operation:

OPEN APPENDECTOMY.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with surrounding lymph nodes

Anesthésie / Anesthesia: General anesthesia with nitrous oxide

Historique et constatations opératoires / History and operative findings:

A 9-year-old non-binary with 1 day history of abdominal pain with anorexia. Failed conservative management for mesenteric adenitis. Imaging: MRI demonstrating RLQ inflammation.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General endotracheal anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. Incision is made in left lower quadrant for open conversion. Accessory port placed in epigastric region. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, extensive adhesions noted. Appendix appeared suppurative, surrounded by localized inflammatory reaction. No abscess, but turbid fluid present. No abnormal adhesions found. The appendix is mobilized using a combination of sharp and blunt dissection. The appendiceal mesentery is carefully taken down with harmonic scalpel. We apply an endoscopic stapler to the base of the appendix and divide it. Specimen placed in EndoCatch bag for removal. We irrigate the abdominal cavity copiously with warm saline until the effluent is clear. We close the fascia with Polysorb 2-0 in a interrupted fashion. Skin is approximated with Steri-Strips.

Monitor for signs of infection; advance diet as tolerated. No mesenteric ischemia. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-YMMXZ2-11774