# PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Children's Hospital Date: 2024-02-22 20:35

Anesthésiste / Anesthetist: Dr. Patricia Wong

Chirurgien / Surgeon: Dr. Daniel Fortin Assistant(s): Dr. resident Anna Kim

Diagnostic préopératoire / Pre-operative diagnosis:

### APPENDICITIS WITH INTUSSUSCEPTION.

Diagnostic postopératoire / Post-operative diagnosis:

SUPPURATIVE APPENDICITIS.

Opération / Operation:

#### APPENDECTOMY WITH LYSIS OF ADHESIONS.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with attached omentum

Anesthésie / Anesthesia: General anesthesia with nitrous oxide

# Historique et constatations opératoires / History and operative findings:

Pediatric patient (17, non-binary) presenting with acute onset right lower quadrant pain. History: no prior abdominal surgery. Imaging confirmed appendicitis.

## Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with mask induction administered. Time-out was performed and abdomen prepped in sterile fashion. Incision is made in left lower quadrant for open conversion. Accessory port placed in epigastric region. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, significant adhesions noted. Appendix appeared friable, surrounded by fluctuating inflammatory reaction. A small localized abscess was found and drained. The appendix was adhered to surrounding structures. The appendix is mobilized using a combination of sharp and blunt dissection. The appendiceal mesentery is carefully taken down with harmonic scalpel. The base of the appendix is healthy and we place three EndoLoops - two proximal and one distal - before transecting the appendix. Specimen placed in EndoCatch bag for removal. We irrigate the abdominal cavity copiously with warm saline until the effluent is clear. Fascial closure is performed at the umbilical site using Vicryl 2-0. The skin is closed with Dermabond.

Pain control with acetaminophen and morphine as needed. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-MZ3445-10377