PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Shriners Hospitals for Children Date: 2024-02-27 22:22

Anesthésiste / Anesthetist: Dr. Camille Roy Chirurgien / Surgeon: Dr. Robert Tremblay

Assistant(s): Dr. resident Leo Morel

Diagnostic préopératoire / Pre-operative diagnosis:

PERFORATED APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

PERFORATED APPENDICITIS WITH ABSCESS.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and peri-appendiceal tissue

Anesthésie / Anesthesia: General anesthesia

Historique et constatations opératoires / History and operative findings:

16-year-old female with 1 day abdominal pain, markedly elevated WBC, elevated CRP, high fever. Imaging: ultrasound showing perforated appendicitis. recent antibiotic use. Past medical history is otherwise unremarkable.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. Total intravenous anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. Transverse infraumbilical incision is performed and access gained via blunt dissection. We place two additional trocars, one in the suprapubic region and one in the left lower quadrant, both under direct vision. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, significant adhesions noted. Appendix appeared gangrenous, surrounded by patchy inflammatory reaction. Multiple small abscesses were encountered. Dense adhesions were encountered during dissection. We proceed with careful dissection of the appendiceal attachments. Appendiceal vessels controlled with clips. Absorbable ligatures are applied prior to amputation. Specimen placed in EndoCatch bag for removal. Copious irrigation is undertaken to ensure removal of all inflammatory debris. We close the fascia with Vicryl 2-0 in a interrupted fashion. Skin is approximated with Dermabond.

Monitor for signs of infection; advance diet as tolerated. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-Y7AQ9B-12934