

PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Charles-LeMoyne

Date: 2023-11-27 22:16

Anesthésiste / Anesthetist: Dr. David Smith

Chirurgien / Surgeon: Dr. James Wilson

Assistant(s): Dr. resident Fatima Sheikh

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH SEPSIS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH BOWEL OBSTRUCTION.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH IRRIGATION AND DRAINAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and inflamed tissue

Anesthésie / Anesthesia: General anesthesia with caudal block

Historique et constatations opératoires / History and operative findings:

Patient (10 years, male) presented with diffuse abdominal pain localizing to RLQ, normal WBC, normal CRP. Imaging: ultrasound reporting lymphadenopathy. Past medical history is otherwise unremarkable.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia and epidural block administered. Time-out was performed and abdomen prepped in sterile fashion. We make an infraumbilical incision of 1 cm, dissect the subcutaneous tissue bluntly and penetrate the abdominal cavity via an open technique. Accessory port placed in epigastric region. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, significant adhesions noted. The surgical field demonstrated gangrenous appendix with fluctuating surrounding inflammation. Purulent fluid was noted throughout the abdominal cavity. The appendix was adhered to surrounding structures. No mesenteric ischemia. Appendix is isolated after adhesiolysis. The mesoappendix is divided using a bipolar energy device. The base of the appendix is healthy and we place three EndoLoops - two proximal and one distal - before transecting the appendix. Specimen placed in EndoCatch bag for removal. We lavage the abdomen extensively, paying particular attention to the pelvis and right gutter. All port sites closed with subcuticular Vicryl 4-0.

IV fluids and pain management as per protocol. Small serosal tear repaired intraoperatively. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-TQ64KR-10110