PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Notre-Dame Date: 2023-12-04 19:28

Anesthésiste / Anesthetist: Dr. Michael Brown

Chirurgien / Surgeon: Dr. Olivia Davis Assistant(s): Dr. resident Anna Kim

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH ABSCESS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH LOCALIZED PERITONITIS.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and inflamed tissue

Anesthésie / Anesthesia: Total intravenous anesthesia

Historique et constatations opératoires / History and operative findings:

4-year-old non-binary with several hours abdominal pain, normal WBC, normal CRP, high fever. Imaging: CT scan showing peri-appendiceal fluid. recent antibiotic use.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. Total intravenous anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. We make an infraumbilical incision of 1 cm, dissect the subcutaneous tissue bluntly and penetrate the abdominal cavity via an open technique. Additional 5 mm trocars are placed in the right and left lower quadrants under laparoscopic guidance. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, diffuse adhesions noted. The appendix was shrunken with patchy inflammation. No abscess was identified. Fibrinous adhesions were lysed during the procedure. Meticulous dissection performed due to distorted anatomy. We dissect the mesentery of the appendix and use the electrocautery to coagulate the artery. We apply an endoscopic stapler to the base of the appendix and divide it. Specimen placed in EndoCatch bag for removal. Thorough irrigation of the abdominal cavity is performed, removing all purulent material. The umbilical fascia is reapproximated with interrupted Polysorb 2-0 sutures. Skin incisions are closed with interrupted silk 4-0.

Repeat CBC and CRP postoperatively. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-10A11P-13114