## PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Pierre-Boucher Date: 2025-08-15 05:12

Anesthésiste / Anesthetist: Dr. John Evans Chirurgien / Surgeon: Dr. Victor Chen

Assistant(s): Dr. resident Chloe Nguyen

Diagnostic préopératoire / Pre-operative diagnosis:

## **COMPLICATED APPENDICITIS.**

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH PERITONEAL CONTAMINATION.

Opération / Operation:

#### APPENDECTOMY WITH REMOVAL OF APPENDICOLITH.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and peri-appendiceal tissue

Anesthésie / Anesthesia: General anesthesia with mask induction

# Historique et constatations opératoires / History and operative findings:

17-year-old female with 1 day abdominal pain, normal WBC, normal CRP, low-grade fever. Imaging: ultrasound showing perforated appendicitis. recent antibiotic use. Past medical history is otherwise unremarkable.

# Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. Total intravenous anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. A small infraumbilical incision is made and the abdominal cavity is entered under direct vision. Additional 5 mm trocars are placed in the right and left lower quadrants under laparoscopic guidance. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, extensive adhesions noted. Operative findings included acutely inflamed appendix and mild inflammation. No abscess, but turbid fluid present. Mild adhesions between bowel loops observed. No evidence of peritoneal carcinomatosis. Appendix is isolated after adhesiolysis. Appendiceal vessels controlled with clips. We secure the appendiceal base with two Endoloops and transect between them. Specimen placed in EndoCatch bag for removal. We irrigate the abdominal cavity copiously with warm saline until the effluent is clear. We close the fascia with PDS 2-0 in a interrupted fashion. Skin is approximated with subcuticular Monocryl 4-0.

IV fluids and pain management as per protocol. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-5ED0QY-10267