## PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Maisonneuve-Rosemont Date: 2024-10-31 07:13

Anesthésiste / Anesthetist: Dr. Julia Miller Chirurgien / Surgeon: Dr. James Wilson Assistant(s): Dr. resident Leo Morel

Diagnostic préopératoire / Pre-operative diagnosis:

## APPENDICITIS WITH FREE FLUID.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH FREE FLUID.

Opération / Operation:

#### APPENDECTOMY WITH REMOVAL OF APPENDICOLITH.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and mesoappendix

Anesthésie / Anesthesia: General anesthesia with mask induction

# Historique et constatations opératoires / History and operative findings:

11-year-old non-binary with 2 days abdominal pain, normal WBC, normal CRP, no fever. Imaging: CT scan confirming appendicitis. recent travel.

# Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with mask induction administered. Time-out was performed and abdomen prepped in sterile fashion. A small infraumbilical incision is made and the abdominal cavity is entered under direct vision. Two working ports are established in the right and left lower quadrants. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, diffuse adhesions noted. Appendix appeared perforated, surrounded by extensive inflammatory reaction. There was a contained abscess in the right lower quadrant. Minimal adhesions were noted. Bladder and ureters visualized, no injury. We proceed with careful dissection of the appendiceal attachments. The mesoappendix is dissected and the appendiceal artery is controlled with electrocautery. We secure the appendiceal base with two Endoloops and transect between them. Specimen placed in EndoCatch bag for removal. Irrigation performed with antibiotic solution. Umbilical port site is closed with PDS 3-0 and skin with Dermabond.

Patient will remain on IV antibiotics with gradual diet advancement as per surgery protocol. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-2JL8IR-10858