## PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Pierre-Boucher Date: 2024-09-03 15:10

Anesthésiste / Anesthetist: Dr. John Evans Chirurgien / Surgeon: Dr. Paul Lambert Assistant(s): Dr. resident Leo Morel

Diagnostic préopératoire / Pre-operative diagnosis:

## LOCALIZED PERITONITIS SECONDARY TO APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH HEMORRHAGE.

Opération / Operation:

#### APPENDECTOMY WITH INTRA-ABDOMINAL DRAINAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix

Anesthésie / Anesthesia: Total intravenous anesthesia

# Historique et constatations opératoires / History and operative findings:

16-year-old male with several hours abdominal pain, normal WBC, normal CRP, no fever. Imaging: ultrasound reporting lymphadenopathy. family history of appendicitis.

## Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with caudal block administered. Time-out was performed and abdomen prepped in sterile fashion. Direct trocar insertion after skin and fascia incision. Two working ports are established in the right and left lower quadrants. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, extensive adhesions noted. Intraoperative examination revealed gangrenous appendix. A large pelvic abscess was present and evacuated. Fibrinous adhesions were lysed during the procedure. The surrounding tissues showed intense reaction. The appendix is mobilized using a combination of sharp and blunt dissection. The mesoappendix is divided using a bipolar energy device. We apply an endoscopic stapler to the base of the appendix and divide it. Specimen placed in EndoCatch bag for removal. Copious irrigation is undertaken to ensure removal of all inflammatory debris. We close the fascia with PDS 3-0 in a interrupted fashion. Skin is approximated with subcuticular Monocryl 4-0.

Pain control with acetaminophen and morphine as needed. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-RY09SI-10202