PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: CHU Sainte-Justine Date: 2025-08-16 17:08

Anesthésiste / Anesthetist: Dr. Paul Anderson Chirurgien / Surgeon: Dr. Martin Levesque Assistant(s): Dr. resident Emily Clark

Diagnostic préopératoire / Pre-operative diagnosis:

ACUTE APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH HEMORRHAGE.

Opération / Operation:

APPENDECTOMY WITH LYSIS OF ADHESIONS.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and mesoappendix

Anesthésie / Anesthesia: General endotracheal anesthesia

Historique et constatations opératoires / History and operative findings:

10-year-old female with one week abdominal pain, normal WBC, high CRP, low-grade fever. Imaging: CT scan confirming appendicitis. history of constipation.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with mask induction administered. Time-out was performed and abdomen prepped in sterile fashion. A vertical infraumbilical incision is made and carried down to the fascia which is incised sharply. Two working ports are established in the right and left lower quadrants. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, extensive adhesions noted. Operative findings included phlegmonous appendix and extensive inflammation. Purulent fluid was noted throughout the abdominal cavity. Fibrinous adhesions were lysed during the procedure. Minor bleeding controlled with cautery. Blunt dissection is used to free the appendix from surrounding structures. We dissect the mesentery of the appendix and use the electrocautery to coagulate the artery. The appendix is ligated at its base with two absorbable sutures and then amputated. Specimen placed in EndoCatch bag for removal. Saline is used for thorough irrigation of all quadrants. Fascial closure is performed at the umbilical site using Vicryl 3-0. The skin is closed with interrupted nylon 4-0.

Continue surgical care with antibiotic therapy and diet advancement as tolerated. Bladder and ureters visualized, no injury. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-DS3KTS-11517