

PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Ste-Agathe Hospital

Date: 2024-03-08 01:32

Anesthésiste / Anesthetist: Dr. Michael Brown

Chirurgien / Surgeon: Dr. Marie-Claire Dubois

Assistant(s): Dr. resident Jennifer Park

Diagnostic préopératoire / Pre-operative diagnosis:

PERFORATED APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

PHLEGMONOUS APPENDICITIS.

Opération / Operation:

APPENDECTOMY WITH LYSIS OF ADHESIONS.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendicolith

Anesthésie / Anesthesia: General anesthesia and epidural block

Historique et constatations opératoires / History and operative findings:

Patient (14 years, male) presented with abdominal pain after trauma, elevated WBC, elevated CRP. Imaging: CT scan showing appendiceal abscess. Past medical history is otherwise unremarkable.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with mask induction administered. Time-out was performed and abdomen prepped in sterile fashion. Transverse infraumbilical incision is performed and access gained via blunt dissection. Three trocars in total are used for laparoscopic access. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, significant adhesions noted. Intraoperative examination revealed perforated appendix. Multiple small abscesses were encountered. Severe adhesions required careful lysis. The surrounding tissues showed fluctuating reaction. No need for drains postoperatively. We carefully dissect the inflammatory mass and identify the perforated appendix. The mesoappendix is divided using a bipolar energy device. The base of the appendix is healthy and we place three EndoLoops - two proximal and one distal - before transecting the appendix. Specimen placed in EndoCatch bag for removal. Irrigation performed with antibiotic solution. We close the fascia with PDS 3-0 in a interrupted fashion. Skin is approximated with Steri-Strips. No unexpected findings.

Continue surgical care with antibiotic therapy and diet advancement as tolerated. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-KDRSPL-13313