PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Jewish General Hospital Date: 2024-05-18 13:23

Anesthésiste / Anesthetist: Dr. Rachel Stein Chirurgien / Surgeon: Dr. Paul Lambert Assistant(s): Dr. resident Jennifer Park

Diagnostic préopératoire / Pre-operative diagnosis:

ACUTE APPENDICITIS WITH PERITONITIS.

Diagnostic postopératoire / Post-operative diagnosis:

ACUTE APPENDICITIS WITH PERITONITIS.

Opération / Operation:

LAPAROSCOPIC CONVERTED TO OPEN APPENDECTOMY.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and inflamed tissue

Anesthésie / Anesthesia: General anesthesia with caudal block

Historique et constatations opératoires / History and operative findings:

10-year-old non-binary with 1 day abdominal pain. Treated for mesenteric adenitis; symptoms persisted. Imaging: MRI demonstrating RLQ inflammation.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with local infiltration administered. Time-out was performed and abdomen prepped in sterile fashion. We create a 1 cm infraumbilical incision and enter the abdomen using the Hasson technique. Supplementary ports are placed in the suprapubic area and left iliac fossa. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, extensive adhesions noted. Findings include hyperemic appendix with intense inflammatory changes. A large pelvic abscess was present and evacuated. No abnormal adhesions found. Appendix is isolated after adhesiolysis. The appendiceal mesentery is carefully taken down with harmonic scalpel. The appendix is ligated at its base with two absorbable sutures and then amputated. Specimen placed in EndoCatch bag for removal. Abdominal lavage performed until clear. We close the fascia with Maxon 2-0 in a interrupted fashion. Skin is approximated with subcuticular Monocryl 4-0. Ovaries and uterus normal in female patients.

Patient to receive postoperative IV antibiotics with monitoring for return of bowel function. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-52IYHD-12643