PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Notre-Dame Date: 2024-05-08 23:49

Anesthésiste / Anesthetist: Dr. John Evans Chirurgien / Surgeon: Dr. Samuel Lee

Assistant(s): Dr. resident Marc Gagnon

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH SEPSIS.

Diagnostic postopératoire / Post-operative diagnosis:

SUPPURATIVE APPENDICITIS.

Opération / Operation:

APPENDECTOMY WITH INTRA-ABDOMINAL DRAINAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and inflamed tissue

Anesthésie / Anesthesia: General anesthesia

Historique et constatations opératoires / History and operative findings:

Pediatric patient (1, non-binary) presenting with acute onset right lower quadrant pain. History: family history of appendicitis. Imaging confirmed appendicitis.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia and epidural block administered. Time-out was performed and abdomen prepped in sterile fashion. Direct trocar insertion after skin and fascia incision. Supplementary ports are placed in the suprapubic area and left iliac fossa. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, moderate adhesions noted. The surgical field demonstrated sclerotic appendix with severe surrounding inflammation. Multiple small abscesses were encountered. Multiple bowel loops adherent to the mass. No evidence of malignancy. We proceed with careful dissection of the appendiceal attachments. The mesoappendix is divided using a bipolar energy device. Endoscopic stapling is used for appendiceal division. Specimen placed in EndoCatch bag for removal. We lavage the abdomen extensively, paying particular attention to the pelvis and right gutter. The umbilical fascia is reapproximated with interrupted Vicryl 2-0 sutures. Skin incisions are closed with interrupted silk 4-0.

Patient will remain on IV antibiotics with gradual diet advancement as per surgery protocol. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-IP53AC-10176