PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Montreal Children's Date: 2025-04-29 11:51

Anesthésiste / Anesthetist: Dr. Paul Anderson Chirurgien / Surgeon: Dr. Elena Rodriguez Assistant(s): Dr. resident Fatima Sheikh

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH FREE FLUID.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH PERITONEAL CONTAMINATION.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH OMENTAL WRAPPING.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix Anesthésie / Anesthesia: General anesthesia with regional block

Historique et constatations opératoires / History and operative findings:

Pediatric patient (11, male) presenting with acute onset abdominal pain after trauma. History: recent antibiotic use. Imaging confirmed appendicitis.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with caudal block administered. Time-out was performed and abdomen prepped in sterile fashion. A vertical infraumbilical incision is made and carried down to the fascia which is incised sharply. Two working ports are established in the right and left lower quadrants. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, minimal adhesions noted. The surgical field demonstrated distended appendix with persistent surrounding inflammation. No abscess was identified. The omentum was wrapped around the inflamed appendix. Minimal intraoperative blood loss. The appendix is mobilized using a combination of sharp and blunt dissection. The appendiceal mesentery is carefully taken down with harmonic scalpel. We apply an endoscopic stapler to the base of the appendix and divide it. Specimen placed in EndoCatch bag for removal. Saline is used for thorough irrigation of all quadrants. The umbilical fascia is reapproximated with interrupted Ethibond 2-0 sutures. Skin incisions are closed with Steri-Strips.

IV fluids and pain management as per protocol. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-6PBO1W-12971