PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: St. Mary's Hospital Date: 2024-07-26 00:15

Anesthésiste / Anesthetist: Dr. John Evans Chirurgien / Surgeon: Dr. Samuel Lee Assistant(s): Dr. resident Emily Clark

Diagnostic préopératoire / Pre-operative diagnosis:

ACUTE APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH PERITONEAL CONTAMINATION.

Opération / Operation:

OPEN APPENDECTOMY WITH PELVIC LAVAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and peri-appendiceal tissue Anesthésie / Anesthesia: General anesthesia with mask induction

Historique et constatations opératoires / History and operative findings:

15-year-old non-binary with one week abdominal pain. Treated for gastroenteritis; symptoms persisted. Imaging: CT scan confirming appendicitis.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia and epidural block administered. Time-out was performed and abdomen prepped in sterile fashion. Transverse infraumbilical incision is performed and access gained via blunt dissection. We place two additional trocars, one in the suprapubic region and one in the left lower quadrant, both under direct vision. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, significant adhesions noted. Appendix appeared hyperemic, surrounded by mild inflammatory reaction. Multiple small abscesses were encountered. Fibrinous adhesions were lysed during the procedure. The appendix is mobilized using a combination of sharp and blunt dissection. The appendiceal artery is ligated and divided. We secure the appendiceal base with two Endoloops and transect between them. Specimen placed in EndoCatch bag for removal. We lavage the abdomen extensively, paying particular attention to the pelvis and right gutter. We close the fascia with Vicryl 3-0 in a interrupted fashion. Skin is approximated with subcuticular Vicryl 4-0.

Pain control with acetaminophen and morphine as needed. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-YPXU5S-12977