# PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Jewish General Hospital Date: 2024-02-14 08:18

Anesthésiste / Anesthetist: Dr. Amélie Moreau

Chirurgien / Surgeon: Dr. Ahmed Khan Assistant(s): Dr. resident Leo Morel

Diagnostic préopératoire / Pre-operative diagnosis:

# APPENDICITIS WITH FREE FLUID.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH HEMORRHAGE.

Opération / Operation:

### LAPAROSCOPIC APPENDECTOMY WITH DRAINAGE OF ABSCESS.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and mesoappendix

Anesthésie / Anesthesia: General anesthesia

#### Historique et constatations opératoires / History and operative findings:

17-year-old male with 2 days abdominal pain. Treated for viral syndrome; symptoms persisted. Imaging: ultrasound suggestive of appendicitis.

# Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. Total intravenous anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. We make an infraumbilical incision of 1 cm, dissect the subcutaneous tissue bluntly and penetrate the abdominal cavity via an open technique. Two working ports are established in the right and left lower quadrants. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, minimal adhesions noted. Operative findings included distended appendix and intense inflammation. No pus or abscess formation found. Minimal adhesions were noted. Minor bleeding controlled with cautery. Blunt dissection is used to free the appendix from surrounding structures. The mesoappendix is divided using a bipolar energy device. We apply an endoscopic stapler to the base of the appendix and divide it. Specimen placed in EndoCatch bag for removal. Copious irrigation is undertaken to ensure removal of all inflammatory debris. We close the fascia with Vicryl 3-0 in a interrupted fashion. Skin is approximated with subcuticular Monocryl 4-0.

IV fluids and pain management as per protocol. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-RZTNZA-10596