## PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Charles-LeMoyne Date: 2025-06-17 07:48

Anesthésiste / Anesthetist: Dr. Rachel Stein

Chirurgien / Surgeon: Dr. Victor Chen Assistant(s): Dr. resident Ethan Wright

Diagnostic préopératoire / Pre-operative diagnosis:

SUPPURATIVE APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

SUPPURATIVE APPENDICITIS.

Opération / Operation:

## LAPAROSCOPIC APPENDECTOMY WITH IRRIGATION AND DRAINAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with attached omentum

Anesthésie / Anesthesia: Total intravenous anesthesia

## Historique et constatations opératoires / History and operative findings:

Pediatric patient (4, non-binary) presenting with acute onset abdominal pain and constipation. History: recent antibiotic use. Imaging confirmed appendicitis. Recent travel history may be relevant.

## Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with nitrous oxide administered. Time-out was performed and abdomen prepped in sterile fashion. Transverse infraumbilical incision is performed and access gained via blunt dissection. Supplementary ports are placed in the suprapubic area and left iliac fossa. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, extensive adhesions noted. Findings include thick-walled appendix with persistent inflammatory changes. No pus or abscess formation found. No significant adhesions. We proceed with careful dissection of the appendiceal attachments. We dissect the mesentery of the appendix and use the electrocautery to coagulate the artery. Base of appendix secured with purse-string suture prior to removal. Specimen placed in EndoCatch bag for removal. Thorough irrigation of the abdominal cavity is performed, removing all purulent material. Fascial closure is performed at the umbilical site using Vicryl 2-0. The skin is closed with interrupted silk 4-0.

Repeat CBC and CRP postoperatively. Patient tolerated procedure well. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-GKKFIC-11273