PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: CHU Sainte-Justine Date: 2025-05-04 18:36

Anesthésiste / Anesthetist: Dr. Claire Dubois Chirurgien / Surgeon: Dr. Martin Levesque Assistant(s): Dr. resident Ethan Wright

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH SEPSIS.

Diagnostic postopératoire / Post-operative diagnosis:

SUPPURATIVE APPENDICITIS.

Opération / Operation:

APPENDECTOMY WITH LYSIS OF ADHESIONS.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and mesoappendix

Anesthésie / Anesthesia: General anesthesia with local infiltration

Historique et constatations opératoires / History and operative findings:

9-year-old female with several hours abdominal pain, markedly elevated WBC, high CRP, no fever. Imaging: ultrasound showing appendicitis. history of constipation.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. Total intravenous anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. Direct trocar insertion after skin and fascia incision. Supplementary ports are placed in the suprapubic area and left iliac fossa. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, diffuse adhesions noted. Appendix was acutely inflamed and surrounded by Abscess cavity found in RLQ and irrigated. and Minimal adhesions were noted. No need for drains postoperatively. Blunt dissection is used to free the appendix from surrounding structures. Mesenteric vessels to the appendix are secured prior to removal. The appendix is ligated at its base with two absorbable sutures and then amputated. Specimen placed in EndoCatch bag for removal. Copious irrigation is undertaken to ensure removal of all inflammatory debris. We close the fascia at the umbilical port with figure-of-eight sutures of Polysorb 2-0 and the skin with subcuticular Vicryl 4-0. No evidence of malignancy.

Patient will remain on IV antibiotics with gradual diet advancement as per surgery protocol. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-T680MU-12936