

PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: St. Mary's Hospital

Date: 2025-05-21 08:15

Anesthésiste / Anesthetist: Dr. Michael Brown

Chirurgien / Surgeon: Dr. Aisha Patel

Assistant(s): Dr. resident Anna Kim

Diagnostic préopératoire / Pre-operative diagnosis:

COMPLICATED APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

PERFORATED APPENDICITIS WITH ABSCESS.

Opération / Operation:

APPENDECTOMY WITH REMOVAL OF APPENDICOLITH.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with surrounding lymph nodes

Anesthésie / Anesthesia: General anesthesia with local infiltration

Historique et constatations opératoires / History and operative findings:

16-year-old female with several hours abdominal pain, normal WBC, elevated CRP, low-grade fever. Imaging: ultrasound suggestive of appendicitis. recent travel. Recent travel history may be relevant.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with local infiltration administered. Time-out was performed and abdomen prepped in sterile fashion. We make an infraumbilical incision of 1 cm, dissect the subcutaneous tissue bluntly and penetrate the abdominal cavity via an open technique. Additional 5 mm trocars are placed in the right and left lower quadrants under laparoscopic guidance. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, minimal adhesions noted. Intraoperative examination revealed sclerotic appendix. A small localized abscess was found and drained. No significant adhesions. The surrounding tissues showed persistent reaction. Blunt dissection is used to free the appendix from surrounding structures. Mesenteric vessels to the appendix are secured prior to removal. The appendix is ligated at its base with two absorbable sutures and then amputated. Specimen placed in EndoCatch bag for removal. Copious irrigation is undertaken to ensure removal of all inflammatory debris. We close the fascia at the umbilical port with figure-of-eight sutures of Maxon 2-0 and the skin with subcuticular Monocryl 4-0.

Repeat CBC and CRP postoperatively. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-6QX84U-11894