PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Notre-Dame Date: 2024-11-02 22:36

Anesthésiste / Anesthetist: Dr. Omar Fahmy

Chirurgien / Surgeon: Dr. Sophie Chen Assistant(s): Dr. resident Zoe Tremblay

Diagnostic préopératoire / Pre-operative diagnosis:

LOCALIZED PERITONITIS SECONDARY TO APPENDICITIS.

Diagnostic postopératoire / Post-operative diagnosis:

ACUTE APPENDICITIS.

Opération / Operation:

APPENDECTOMY WITH REMOVAL OF APPENDICOLITH.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and mesoappendix

Anesthésie / Anesthesia: General endotracheal anesthesia

Historique et constatations opératoires / History and operative findings:

A 5-year-old male who presented with abdominal pain with rebound tenderness. Initially evaluated 3 days prior and diagnosed with ovarian cyst. Now has elevated WBC, normal CRP, no fever. Imaging: MRI demonstrating RLQ inflammation. Past medical history is otherwise unremarkable.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia with nitrous oxide administered. Time-out was performed and abdomen prepped in sterile fashion. We make an infraumbilical incision of 1 cm, dissect the subcutaneous tissue bluntly and penetrate the abdominal cavity via an open technique. Accessory port placed in epigastric region. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, extensive adhesions noted. Findings include friable appendix with moderate inflammatory changes. No abscess was identified. Fibrinous adhesions were lysed during the procedure. No evidence of malignancy. We carefully dissect the inflammatory mass and identify the thick-walled appendix. The appendiceal mesentery is carefully taken down with harmonic scalpel. Endoscopic stapling is used for appendiceal division. Specimen placed in EndoCatch bag for removal. Copious irrigation is undertaken to ensure removal of all inflammatory debris. All port sites closed with non-absorbable Prolene 4-0.

IV fluids and pain management as per protocol. Small serosal tear repaired intraoperatively. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-A10GH9-13462